

What You Need to Know About CEFIXIME

This medication is used to cure you of an infection.
Contact your healthcare provider, if your symptoms do not improve or get worse.

HOW TO TAKE CEFIXIME:

Take two tablets by mouth as a single dose

CAUTION

- ✔ Tell your healthcare provider if you have any kidney or liver diseases, bleeding disorders, or disorders that result in the build up of diet amino acid
- ✔ If you are taking any prescription, non-prescription, herbal, or recreational products, please discuss with your healthcare provider
- ✔ Use a condom, as this medication can interact with hormonal birth control and make them less effective

POSSIBLE SIDE EFFECTS

- You may experience mild diarrhea, abdominal pain or nausea
- If any of these effects persist or worsen, please contact your healthcare provider

AFTER TAKING THIS MEDICATION

- ✔ Do not have sex for one week after you and your partner(s) have completed treatment and all symptoms have resolved
- ✔ Tell your healthcare provider if symptoms do not improve or resolve in 5 days
- ✔ Drink plenty of water
- ✔ Cefixime may interact with certain urine glucose or ketone tests
- ✔ Cefixime may affect the result of certain lab tests



ALERT your healthcare provider if you have any of the issues below:

- 🚫 Diarrhea that is watery or bloody
- 🚫 Are pregnant or may become pregnant
- 🚫 Develop a skin rash, have trouble breathing, or notice swelling in your face or throat
- 🚫 Worsening of joint pain, fever, or enlarged lymph nodes
- 🚫 Have a seizure



For more information, please scan this QR code:



Rev Nov 2022

