What You Need to Know About DOXYCYCLINE

This medication is used to cure an infection.

It is important to <u>take the entire prescription even if you are not having symptoms or your symptoms have resolved</u> to treat your infection and protect your partner(s).

HOW TO TAKE DOXYCYCLINE:

Take one capsule by mouth twice a day for seven days

WHILE ON THIS MEDICATION

- Do not have sex while taking this medication, and until all symptoms have resolved to prevent spreading the infection to your partner(s)
- Use a condom as this medication can interact with hormonal birth control and make them less effective
- Do not take supplements containing iron or calcium, multivitamins, antacids, or laxatives within 2 hours before or after taking doxycycline
- Drink plenty of water
- ◆ Cover up when going out in the sun and wear sunscreen (SPF 30+)— this medication can make you sunburn more easily
- Tell your health care provider if symptoms do not improve or resolve in 5 days

POSSIBLE SIDE EFFECTS

- You may experience mild diarrhea, upset stomach, or nausea
- If any of these effects persist or worsen, please contact your healthcare provider

TIPS TO HELP YOU TAKE DOXYCYCLINE

- ◆ Take this medication with a full glass of water and do not lay down for 30 minutes after to avoid throat irritation
- Take this medication with food to help with upset stomach
- ◆ Place a reminder note on an object or set an alarm on your phone to remind you to take your medication at the same time every day
- Mark off on a calendar that you have taken your medication.

IF YOU FORGET TO TAKE A DOSE

- Take the missed dose as soon as you remember
 - **UNLESS** it is almost time for your next dose, **THEN SKIP** the missed dose and take the next one at the scheduled time
- DO NOT take any extra capsules to make up for the missed dose



STOP taking your medicine right away if you have any of the issues below:

- Diarrhea that is watery or bloody
- Are pregnant or may become pregnant
- Develop a skin rash, have trouble breathing, or notice swelling in your face or throat
- Develop rapid headache, blurred vision, double vision, or vision loss



For more information, please scan this QR code:



